



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07C 209/42, 211/52	A1	(11) International Publication Number: WO 00/69805 (43) International Publication Date: 23 November 2000 (23.11.00)
(21) International Application Number: PCT/EP00/04595 (22) International Filing Date: 11 May 2000 (11.05.00) (30) Priority Data: 9911180.9 13 May 1999 (13.05.99) GB (71) Applicant (for all designated States except US): AVENTIS CROPSCIENCE S.A. [FR/FR]; 55, avenue René Cassin, F-69009 Lyon (FR). (72) Inventors; and (75) Inventors/Applicants (for US only): ANCEL, Jean-Erick [FR/FR]; 14, rue Lucien Begule, F-69230 Saint-Genis-Laval (FR). PERRIN-JANET, Gilles [FR/FR]; 17, allée des Clementières, F-69270 Chaponnay (FR). LEROY, Pierre [FR/FR]; 21, rue de la Bannière, F-69003 Lyon (FR). (74) Agent: BRACHOTTE, Charles; Aventis CropScience S.A., Département Propriété Industrielle, B.P. 9163, F-69263 Lyon Cedex 09 (FR).	(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: PROCESSES FOR PREPARING PESTICIDAL INTERMEDIATES		
<div style="text-align: center;"> (1)</div>		
(57) Abstract <p>The invention relates to processes for the preparation of compounds of formula (I), wherein R¹ and R² each independently represent a halogen atom.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Processes For Preparing
Pesticidal Intermediates


This invention relates to novel processes for preparing intermediates (particularly certain aniline compounds and phenylhydrazine derivatives) useful in the preparation of pesticides.

3,5-Dihaloanilines and particularly 3,5-dichloroaniline are important synthetic intermediates for a number of valuable compounds including agrochemical fungicides for example as described in US Patent Publication Numbers 3755350 and 3903090, or German Patent Publication No.2906574.

Various methods for preparing these compounds (particularly 3,5-dichloroaniline) are known, for example as described in Japanese patent publication numbers 05194330 and 60048500, and US patent publication number 4508922. However these procedures are generally inefficient, slow and suffer from poor selectivity and low yields. Furthermore they generally require the use of ammonia at high temperatures and/or pressures. There remains a need to develop new methods for obtaining these compounds particularly to find procedures which are safe to operate, efficient and can be performed at moderate temperatures and pressures.


The present applicants have surprisingly discovered novel processes for the preparation of certain substituted anilines and phenylhydrazines which do not suffer from these disadvantages, thus providing a new method for preparing valuable synthetic intermediates.

The present invention accordingly provides a process (A) for the preparation of a substituted aniline of formula (I):



(I)

wherein R¹ and R² each independently represent a halogen atom; or an acid addition salt thereof; which process comprises the hydrogenolysis of a substituted phenylhydrazine of formula (II):



(II)

wherein R¹ and R² are as hereinbefore defined, or an acid addition salt thereof, in the presence of a metal or metal compound (for example a metal salt) under reducing conditions.

Unless otherwise specified in the present specification halogen atoms are selected from fluorine, chlorine, bromine and iodine.

The acid addition salts referred to in the invention are preferably the salts formed from strong acids such as mineral acids, for example sulphuric acid or hydrochloric acid.

The hydrogenolysis may be performed using a metal or metal salt selected from Raney nickel (a nickel-aluminium alloy) optionally in the presence of iron, manganese, cobalt, copper, zinc or chromium; stannous chloride; zinc in the presence of acetic acid; and a molybdenum (III) salt. The reaction may also be carried out using Raney nickel, platinum or palladium (which may be supported on charcoal or other inert material) in the presence of hydrogen gas. When the reaction is performed with hydrogen gas a pressure of 2 to 20 bars (preferably 5 to 10 bars) is generally used. The hydrogenolysis is preferably performed using Raney nickel.


The reaction may be performed in the absence or presence of a solvent, but is generally conducted in a solvent which may be selected from alcohols such as methanol, ethanol, propanol, isopropanol, butanol, isobutanol or tertiary butanol; ethers; aromatic hydrocarbons and water (or mixtures thereof). Methanol and ethanol are preferred solvents.

The reaction temperature is generally from 20°C to 150°C, preferably from 20°C to 90°C. The amount of catalyst employed is generally from 0.01 to 3 molar equivalents (preferably from 0.05 to 2 molar equivalents), although when the reaction is carried out under an atmosphere of hydrogen, a smaller amount generally gives satisfactory results.

Process (A) seeks to enable substituted aniline compounds of formula (I) and their acid addition salts thereof to be obtained in high yield from readily available starting materials. Furthermore the reaction

can be very simple and economical to perform, and product isolation can be straightforward. Another advantage of this method is that these compounds of formula (I) may be prepared at moderate temperatures and pressures, whereas prior art methods require high temperatures and pressures.

According to a further feature of the present invention there is provided a process (B) for the preparation of a substituted phenylhydrazine of formula (II) as defined above, or an acid addition salt thereof, which comprises the reaction of a compound of formula (III):



(III)

wherein R^1 and R^2 are as hereinbefore defined and X represents a halogen atom, with hydrazine or an acid addition salt or source thereof.

By appropriate selection of the halogen atoms which represent R^1 , R^2 and X in formula (III) so that the group X is more reactive than R^1 and R^2 , it is possible to obtain substituted phenylhydrazines of formula (II) and salts thereof having the various combinations of halogen atoms. Particularly preferred compounds are those wherein R^1 and R^2 in formulae (I) and (II) each represent a chlorine atom.

Compounds of formula (III) are generally known.

Preferably hydrazine hydrate is used in the process.

When an acid addition salt of hydrazine is employed a base such as a trialkylamine (for example triethylamine) is optionally present.

A particularly preferred compound of formula (III) is 1,3,5-trichlorobenzene.

The reaction may be conducted in a solvent chosen from cyclic and aliphatic ethers such as tetrahydrofuran, 1,4-dioxan or 1,2-dimethoxyethane; N-methylpyrrolidone; dimethyl sulphoxide; N,N-dimethylformamide; sulfolane; N,N,N',N'-tetramethylurea; aromatic hydrocarbons which may be substituted with one or more alkyl groups or chlorine atoms, such as chlorobenzene or xylene; and pyridine.

Preferred solvents include pyridine, tetrahydrofuran,

N,N,N',N'-tetramethylurea and 1,4-dioxan (pyridine and tetrahydrofuran

are especially preferred). The amount of solvent used is generally from 1 to 10ml (preferably from 4 to 8ml) per gramme of compound of formula (III).

The process is generally performed in an autoclave or other sealed vessel.

The reaction temperature is generally from 50°C to 250°C, preferably from 120°C to 200°C. The reaction is generally conducted using from 1 to 20 molar equivalents (preferably 4 to 8 molar equivalents) of the hydrazine source.

A catalyst may optionally be used to increase the rate of the reaction, and when present is generally chosen from alkali and alkaline earth metal fluorides such as potassium fluoride. The amount of catalyst employed is generally from 0.05 to 2 molar equivalents (preferably from 0.5 to 1 molar equivalents). The reaction may also be effected in the presence of copper or a copper salt, preferably copper (I) chloride.

The process of the invention for the preparation of substituted phenylhydrazines of formula (II) and their acid addition salts represents in certain aspects an improvement over the prior art. Process (B) is particularly useful for the preparation of the valuable intermediate 3,5-dichlorophenylhydrazine from 1,3,5-trichlorobenzene, because the reaction proceeds in high yield and isomer purity and provides together with the other processes of the present invention an efficient method for obtaining synthetically valuable intermediate compounds.

Moreover when process (B) is used for the preparation of 3,5-dichlorophenylhydrazine from 1,3,5-trichlorobenzene, the reaction proceeds with excellent regioselectivity.

According to a further feature of the invention processes (A) and (B) can be combined to prepare a substituted aniline of formula (I) or an acid addition salt thereof from a compound of formula (III).

The following non-limiting examples illustrate the invention. Each product was shown to be identical to a known reference sample of the compound.

Example 1

Preparation of 3,5-dichloroaniline from 3,5-dichlorophenylhydrazine

Raney nickel (2g of a slurry in water) was added to a solution of 3,5-dichlorophenylhydrazine (1g) in methanol (5ml) and heated at 60°C for 2 hours. The cooled mixture was filtered and evaporated to give the title compound in 100% yield.

5

Example 2**Preparation 3,5-dichloroaniline from 3,5-dichlorophenylhydrazine**

The procedure of Example 1 was repeated but using two equivalents of Raney nickel and heating at 60°C for 3 hours in methanol to give the title compound in >90% yield.

10

Example 3**Preparation of 3,5-dichlorophenylhydrazine**

A mixture of 1,3,5-trichlorobenzene (2.53g), hydrazine hydrate (3.48g, 5 molar equivalents) and pyridine (12ml) was heated in an autoclave (purged with argon) for 6 hours at 180°C. The mixture was cooled, the excess hydrazine decanted and the organic phase evaporated in vacuo to give the title compound in 60% yield. It was shown that 65% of the starting material had been consumed, thus indicating that the reaction had occurred with high selectivity.

15

20

Example 4**Preparation of 3,5-dichlorophenylhydrazine**

The procedure of Example 3 was repeated but using 6 equivalents of hydrazine hydrate gave the title compound in 63% yield. It was shown that 77% of the starting material had been consumed, thus indicating that the reaction had occurred with good selectivity.

25

Example 5**Preparation of 3,5-dichlorophenylhydrazine**

The procedure of Example 3 was repeated but using 6 equivalents of hydrazine hydrate to give after 10 hours at 180°C, the title compound in 76% yield. It was shown that 77% of the starting material had been consumed, thus indicating that the reaction had occurred with high selectivity.

30


Example 6 Preparation of 3,5-dichlorophenylhydrazine

The procedure of Example 3 was repeated but using 6 equivalents of hydrazine hydrate to give after 6 hours at 200°C, the title compound in 82% yield. It was shown that 83% of the starting material had been consumed, thus indicating that the reaction had occurred with high selectivity.

35


CLAIMS

1. A process for the preparation of a substituted aniline of formula (I):



(I)

wherein R^1 and R^2 each independently represent a halogen atom; or an acid addition salt thereof; which process comprises the hydrogenolysis of a substituted phenylhydrazine of formula (II):




(II)

wherein R^1 and R^2 are as defined above, or an acid addition salt thereof, in the presence of a metal or metal compound under reducing conditions.

2. A process according to claim 1 in which the hydrogenolysis is performed using Raney nickel.

3. A process for the preparation of a substituted phenylhydrazine of formula (II) as defined in claim 1 or an acid addition salt thereof, which comprises the reaction of a compound of formula (III):



(III)

wherein R^1 and R^2 are as defined in claim 1 and X represents a halogen atom, with hydrazine or an acid addition salt or source thereof.

4. A process according to claim 3 in which hydrazine hydrate is used.

5. A process according to claim 1 or 2 in which the compound of formula (II) is prepared by a process according to claim 3 or 4.

6. A process according to any one of the preceding claims wherein R¹ and R² represent chlorine.

10

7. A process according to claims 1,2,5 or 6 which further comprises converting the substituted aniline of formula (I) or an acid addition salt thereof into an agriculturally fungicidal compound.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/04595

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C209/42 C07C211/52

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, BEILSTEIN Data, WPI Data, PAJ, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	"Methoden der Organischen Chemie (Houben-Weyl) Band E16d", GEORG THIEME VERLAG, STUTTGART-NEW YORK XP002147228 page 985 -page 986	1,2
X	"Methoden der Organischen Chemie (Houben-Weyl) Band E16a" 1990, GEORG THIEME VERLAG, STUTTGART-NEW YORK XP002147227 page 648 -page 653	3-7
A	J.H. WOTIZ ET AL.: "Low temperature amination of aromatic polyhalides" JOURNAL OF ORGANIC CHEMISTRY, vol. 24, 1959, pages 595-598, XP002147225 EASTON US page 597	1
	-/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

27 September 2000

Date of mailing of the international search report

17/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Pauwels, G